


USE OF PROPARGYL GLYCINE AMINO PROPARGYL DIOL COMPOUNDS FOR PREVENTION OF HYPERTENSION

ABSTRACT

5

Compounds characterized generally as propargyl glycine amino propargyl diol derivatives are useful for prevention of hypertension. Compounds of particular interest are those of Formula I

10



wherein A is selected from CO and SO₂ wherein X is selected from oxygen atom and methylene; wherein each of R₁ and R₉ is a group independently selected from hydrido, methyl, ethyl, n-propyl, isopropyl, benzyl, b, b, b-trifluoroethyl, t-butyloxycarbonyl and methoxymethylcarbonyl, and wherein the nitrogen atom to which R₁ and R₉ are attached may be combined with oxygen to form an N-oxide; wherein R₂ is selected from hydrido, methyl, ethyl and isopropyl; wherein R₃ is selected from benzyl, cyclohexylmethyl, phenethyl, imidazolemethyl, pyridylmethyl and 2-pyridylethyl; wherein each of R₅ and R₈ is independently propargyl or a propargyl-containing moiety; wherein R₇ is cyclohexylmethyl; wherein each of R₄ and R₆ is independently selected from hydrido and methyl; wherein each of R₁₁ and R₁₂ is independently selected from hydrido, alkyl and phenyl; wherein m is zero; and wherein n is a number selected from zero through three; or a pharmaceutically-acceptable salt thereof.

 00906343 112004
 T09277 CHE90600